

# Angioedema in chronic spontaneous urticaria is underdiagnosed and has a substantial impact: Analyses from ASSURE-CSU

G. Sussman<sup>1</sup>  | M. Abuzakouk<sup>2</sup>  | F. Bérard<sup>3</sup> | W. Canonica<sup>4</sup>  |  
H. Oude Elberink<sup>5,6</sup> | A. Giménez-Arnau<sup>7</sup>  | C. Grattan<sup>8</sup>  | K. Hollis<sup>9</sup> | S. Hunter<sup>9</sup> |  
A. Knulst<sup>10</sup>  | J.-P. Lacour<sup>11</sup>  | C. Lynde<sup>12</sup> | A. Marsland<sup>13</sup> | D. McBride<sup>14</sup>  |  
M. Maurer<sup>15</sup>  | A. Nakonechna<sup>16</sup>  | J. Ortiz de Frutos<sup>17</sup> | M. Reynolds<sup>9</sup> |  
C. Sweeney<sup>9</sup> | H. Tian<sup>18</sup> | K. Weller<sup>15</sup>  | D. Wolin<sup>9</sup> | M.-M. Balp<sup>19</sup> 

<sup>1</sup>University of Toronto, Toronto, ON, Canada

<sup>2</sup>Cleveland Clinic Abu Dhabi, Abu Dhabi, UAE

<sup>3</sup>Claude Bernard University Lyon, Villeurbanne, France

<sup>4</sup>IRCCS-Humanitas Research Hospital, Humanitas University, Rozzano-Milano, Italy

<sup>5</sup>Department of Allergology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

<sup>6</sup>GRIAC Research Institute, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

<sup>7</sup>Hospital del Mar, Universitat Autònoma, Barcelona, Spain

<sup>8</sup>Guy's Hospital, London, UK

<sup>9</sup>RTI Health Solutions, Research Triangle Park, NC, USA

<sup>10</sup>University Medical Center, Utrecht, The Netherlands

<sup>11</sup>Université Nice Sophia Antipolis, Nice, France

<sup>12</sup>Lynderm Research, Toronto, ON, Canada

<sup>13</sup>Salford Royal Hospital, University of Manchester, Salford, UK

<sup>14</sup>RTI Health Solutions, Manchester, UK

<sup>15</sup>Charité - Universitätsmedizin Berlin, Berlin, Germany

<sup>16</sup>Royal Liverpool and Broadgreen University Hospitals, Liverpool, UK

<sup>17</sup>Hospital 12 Octubre, Madrid, Spain

<sup>18</sup>Novartis Pharmaceuticals Corporation, East Hanover, NJ, USA

<sup>19</sup>Novartis Pharma AG, Basel, Switzerland

## Correspondence

Gordon Sussman, University of Toronto,  
Toronto, ON, Canada.  
Email: gsussman@rogers.com

## Funding information

Novartis Pharma AG

## Abstract

**Background:** ASSURE-CSU revealed differences in physician and patient reporting of angioedema. This post hoc analysis was conducted to evaluate the actual rate of angioedema in the study population and explore differences between patients with and without angioedema.

**Methods:** This international observational study assessed 673 patients with inadequately controlled chronic spontaneous urticaria (CSU). Physicians abstracted

angioedema data from medical records, which were compared with patient-reported data. Patients in the *Yes-angioedema* category had angioedema reported in the medical record and a patient-reported source. For those in the *No-angioedema* category, angioedema was reported in neither the medical record nor a patient-reported source. Those in the *Misaligned* category had angioedema reported in only one source. Statistical comparisons between *Yes-angioedema* and *No-angioedema* categories were conducted for measures of CSU activity, health-related quality of life (HRQoL), productivity and healthcare resource utilization (HCRU). Regression analyses explored the relationship between Dermatology Life Quality Index (DLQI) score and angioedema, adjusting for important covariates.

**Results:** Among evaluable patients, 259 (40.3%), 173 (26.9%) and 211 (32.8%) were in the *Yes-angioedema*, *No-angioedema* and *Misaligned* category, respectively. CSU activity and impact on HRQoL, productivity, and HCRU was greater for *Yes-angioedema* patients than *No-angioedema* patients. After covariate adjustment, mean DLQI score was significantly higher (indicating worse HRQoL) for patients with angioedema versus no angioedema (9.88 vs 7.27,  $P < .001$ ). The *Misaligned* category had similar results with *Yes-angioedema* on all outcomes.

**Conclusions:** Angioedema in CSU seems to be under-reported but has significant negative impacts on HRQoL, daily activities, HCRU and work compared with no angioedema.

#### KEYWORDS

angioedema, economic burden, observational study, quality of life, urticaria

## 1 | INTRODUCTION

Chronic spontaneous urticaria (CSU) is characterized by the presence of hives, angioedema or both, recurring for 6 weeks or longer, in the absence of identifiable triggers.<sup>1,2</sup> Angioedema is defined as rapid swelling at a deeper level under the skin than hives, occurring on the face, inside the mouth or elsewhere on the body.<sup>2,3</sup> It is estimated that 33%-67% of patients with CSU exhibit both hives and angioedema, 29%-65% exhibit only hives and 1%-13% exhibit only angioedema.<sup>4,5</sup>

Patients who experience CSU as concurrent hives and angioedema often experience a longer duration of disease than patients who experience only hives.<sup>6,7</sup> Angioedema may be disfiguring or painful, may limit daily activities and may have a significant impact on quality of life.<sup>8,9</sup> Patients with CSU-associated angioedema often experience concern over their health status, at times worrying that swelling episodes may cause problems with breathing or may be life-threatening.<sup>9</sup> A real-world multicentre study in Germany has shown that in a 6-month period, more than 40% of patients with inadequately controlled CSU experienced angioedema; among these patients, 78% rated their angioedema as severe or moderate in intensity, and a mean of 34 days with angioedema was reported during the 6-month period.<sup>5</sup> In addition to its considerable symptom burden, recurrent angioedema (in those with hereditary angioedema)

#### Highlights

- Nearly one-third of CSU patients in ASSURE-CSU reported having experienced angioedema in the past 12 months but did not have physician-reported angioedema as documented in their medical records.
- Patients with angioedema had greater CSU activity, HRQoL and productivity impairment, and resource utilization than those without. After controlling for other factors, angioedema significantly affected DLQI total score.
- Among patients with inadequately controlled symptomatic CSU, the proportion with angioedema may be higher than previously thought.

can lead to absenteeism and can have a negative impact on work productivity.<sup>10</sup>

The observational, multinational ASSURE-CSU (ASsessment of the Economic and Humanistic Burden of Chronic Spontaneous/Idiopathic Urticaria PatiEnts) study aimed to characterize the patient population with inadequately controlled CSU and to evaluate the burden of disease.<sup>11,12</sup> Analyses of ASSURE-CSU data revealed

different rates of CSU-associated angioedema between physician and patient reports.<sup>12</sup> The objectives of this post hoc analysis were to provide a better estimate of angioedema rates among CSU patients in the ASSURE-CSU study by aligning patient and physician reports and to analyse differences in patient characteristics as well as humanistic and economic burden between individuals with and without angioedema.

## 2 | METHODS

### 2.1 | ASSURE-CSU study design

ASSURE-CSU was an observational, noninterventional, multinational, multicentre study conducted at urticaria specialty centres in Canada, France, Germany, Italy, Spain, the Netherlands and the United Kingdom. Methodology<sup>11</sup> and overall results<sup>12</sup> have been reported previously. In summary, recruitment of 700 patients was planned. Adult patients who had a clinician-confirmed CSU diagnosis, had received at least one treatment course with an H<sub>1</sub>-antihistamine, had been symptomatic for more than 12 months and were currently symptomatic despite treatment were eligible<sup>12</sup>; patients with urticaria that was predominantly of the inducible form were ineligible. Study data were collected via a retrospective 12-month patient medical record abstraction (MRA) by physicians, a cross-sectional patient survey and a prospective 8-day patient diary. The appropriate national-, local- and site-level ethical approvals were obtained, and all patients provided written informed consent. The study complied with the Declaration of Helsinki.<sup>13</sup>

### 2.2 | Study measures

Study measures relevant to this analysis included the occurrence of angioedema as reported in the physician MRA, the cross-sectional patient survey and the patient diary. Specifically, physicians reported from the medical chart whether the patient had angioedema associated with his or her CSU ever, at the time of diagnosis, and/or within the past 12 months. In the patient survey, completed at enrolment, patients answered whether they had experienced angioedema ever, during the past 12 months, within the past 4 weeks and/or currently (at survey completion); patients also answered questions about the duration and location of angioedema, as well as the symptoms they experienced with angioedema (swelling, itching and pain) and what they would do to seek treatment (or not) during an angioedema episode. In the Urticaria Patient Daily Diary (UPDD), completed for each of 7 days following enrolment, patients indicated whether they had experienced "rapid swelling, also called angioedema," during the past 24 hours.

Additional measures explored in this analysis included demographic data, comorbidities and healthcare resource utilization (HCRU) related to CSU during the previous 12 months, as abstracted by physicians in the MRA. Additional measures collected in the patient survey included validated patient-reported outcome measures such as the Chronic Urticaria Quality of Life Questionnaire (CU-Q<sub>2oL</sub>)<sup>14,15</sup> and Dermatology Life Quality Index (DLQI)<sup>16</sup> and

HCRU. The 7-day UPDD<sup>17</sup> also included the twice-daily Urticaria Activity Score over 7 Days (UAS<sub>7TD</sub>)<sup>18</sup>, daily sleep interference and activity impairment, in addition to the aforementioned occurrence and management of angioedema. On the 8th day, patients completed the Work Productivity and Activity Impairment (WPAI), which has a recall period of 7 days.<sup>19</sup>

### 2.3 | Statistical analysis

Frequency of angioedema within the past 12 months and misalignment of angioedema reporting between physician and patient data sources were evaluated. For patients in the *Yes-angioedema* category, physician and patient data sources agreed that the patient had experienced angioedema. For these cases, the MRA and either the patient survey or patient diary indicated that the patient had experienced angioedema in the past 12 months. For patients in the *No-angioedema* category, physician and patient data sources agreed that the patient had not experienced angioedema. For these cases, all 3 data sources indicated no angioedema in the past 12 months. For the *Misaligned* category, the physician and patient data sources (either patient survey or diary) did not agree as to whether the patient had experienced angioedema. For these cases, the physician data source indicated no angioedema during the past 12 months while one of the patient sources indicated that angioedema had occurred, or the physician source indicated that the patient had experienced angioedema in the past 12 months while the patient sources did not. If the angioedema classification was missing either from the MRA or from both patient data sources (survey and diary), the patient was assigned to a Missing category and excluded from analysis.

Descriptive statistics were used to characterize the study population by angioedema classification (*Yes-angioedema*, *No-angioedema*, *Misaligned*). Disease activity was determined by UAS<sub>7TD</sub> score bands: UAS<sub>7TD</sub> = 0-6 (urticaria-free or well-controlled urticaria activity), 7-15 (mild activity), 16-27 (moderate activity) and 28-42 (severe activity).<sup>20</sup> Health-related quality of life (HRQoL) measures were analysed by published methodology where available.

Statistical comparisons of patients with and without angioedema were conducted using *t* tests or Kruskal-Wallis tests for continuous variables and chi-square tests for categorical variables. Tests were performed between those in the *Yes-angioedema* and *No-angioedema* categories; patients who fell into the *Misaligned* category were not included in the statistical comparisons.

Regression modelling was used to explore whether the relationship between DLQI score and angioedema remained significant after adjusting for important covariates. An analysis of covariance model was used to evaluate the effect of angioedema on DLQI total score. Adjustment covariates included UAS<sub>7TD</sub> score, age, sex, country, disease duration from diagnosis to enrolment and selected comorbidities at enrolment (hypersensitivity to nonsteroidal anti-inflammatory drugs, Hashimoto's disease and asthma). UAS<sub>7TD</sub> score, age and disease duration were treated as continuous variables. Least-squares mean estimates and standard errors were estimated for each angioedema classification; patients in the *Misaligned* category were

combined with patients in the *Yes-angioedema* category for the primary analysis. A sensitivity analysis was also conducted in which the *Misaligned* category was excluded from the comparison.

All tests were performed at a nominal significance level of  $\alpha = .05$  with 2-sided, single degree-of-freedom tests. No correction was made for multiple comparisons. All analyses were performed using SAS for Windows statistical software, version 9.4 (SAS Institute, Inc., Cary, NC, USA).

### 3 | RESULTS

#### 3.1 | Initial angioedema frequency over the past 12 months

According to the MRA, physicians reported that 276 of 673 patients enrolled (41.0%) had experienced CSU-associated angioedema within the past 12 months, with a mean (standard deviation [SD]) of 19.0 (42.13) angioedema episodes during this period. Among the 649 patients who completed the survey, 427 (65.8%) patients reported having had angioedema within the past 12 months (Figure 1A).

Among the 614 patients who completed the diary over 1 week, 294 (47.9%) patients reported that they had angioedema at least 1 day; the mean (SD) patient-reported number of days with angioedema was 3.2 (1.92) over the 7 days. The occurrence of angioedema and number of days with angioedema increased with increasing disease activity over that week.<sup>12</sup>

#### 3.2 | Postalignment angioedema frequency over the past 12 months

Of the 673 patients enrolled, 643 patients had recorded data for assigning the angioedema classification, and 30 patients had missing

data either in the MRA or from both patient data sources (survey and diary) and could not be classified. Of the 643 patients included in the angioedema analyses, 276 (42.9%) had angioedema according to the MRA, and 467 (72.6%) had angioedema according to one of the patient sources (survey or diary).


For the analyses, 259 patients (40.3%) were in the *Yes-angioedema* category, 173 (26.9%) were in the *No-angioedema* category, and 211 (32.8%) were in the *Misaligned* category. Of the 211 patients in the *Misaligned* category, 205 (97.2%) had angioedema recorded in one of the patient sources (survey or diary) but not in the MRA (Figure 1B; Table S1).

#### 3.3 | Patient demographics

There were no significant differences in baseline characteristics between patients in the *Yes-angioedema* and *No-angioedema* categories, except in duration of CSU and presence of Hashimoto's disease (Table 1). Patients in the *Yes-angioedema* category had longer mean (SD) duration of disease from diagnosis to enrolment (61.7 [76.64] months) than those in the *No-angioedema* category (46.1 [69.06]) ( $P < .05$ ) and a higher prevalence of Hashimoto's disease (10.4% vs 2.9%;  $P < .01$ ).

#### 3.4 | Urticaria severity and activity

More patients in the *Yes-angioedema* category were reported to have severe CSU activity at diagnosis (37.8%) than those in the *No-angioedema* category (26.0%) (Table 1). Mean (SD) UAS7<sub>TD</sub> score reported in the patient diary was higher for patients in the *Yes-angioedema* category (17.6 [10.55]) than for those in the *No-angioedema* category (14.6 [8.97]) ( $P < .01$ ) (Figure 2). In line with this observation, more patients in the *Yes-angioedema* category than in the *No-angioedema* category



**FIGURE 1** Frequency of angioedema in the past 12 months. (A) Initial frequency: Angioedema in the past 12 months was more frequently reported by patients who completed the survey (65.8%, 427/649 patients) than recorded in the medical record (41.0%, 276/673 patients). (B) Revised frequency: Of the 32.8% of patients (211/643) with misaligned reporting of angioedema in the past 12 months, 97.2% (205/211) reported having had angioedema when it was not noted in the medical record

**TABLE 1** Patient and disease characteristics from the medical record abstraction, overall and by angioedema classification

Patient characteristics <sup>a</sup>	Total (n = 643)	Angioedema category			P value <sup>b</sup> (yes vs no)
		Yes (n = 259)	No (n = 173)	Misaligned (n = 211)	
Age at enrolment (years)					.0906
Mean (SD)	49.0 (15.56)	49.4 (14.45)	46.8 (16.50)	50.4 (15.95)	
Min, Max	19.0, 89.0	20.0, 81.0	19.0, 89.0	19.0, 87.0	
Female sex (%)	469 (72.9)	197 (76.1)	117 (67.6)	155 (73.5)	.0540
Race and ethnicity <sup>c</sup>	544	205	155	184	.5262
Caucasian/White (%)	491 (90.3)	189 (92.2)	138 (89.0)	164 (89.1)	
Other (%)	44 (8.1)	15 (7.3)	14 (9.0)	15 (8.2)	
Data not available (%)	9 (1.7)	1 (0.5)	3 (1.9)	5 (2.7)	
Disease duration from diagnosis to enrolment (months)					.0486
Mean (SD)	57.3 (77.73)	61.7 (76.64)	46.1 (69.06)	61.0 (85.43)	
Comorbid conditions at enrolment (%)					
Hypersensitivity to NSAIDs	52 (8.1)	28 (10.8)	10 (5.8)	14 (6.6)	.0833
Hashimoto's disease	43 (6.7)	27 (10.4)	5 (2.9)	11 (5.2)	.0042
Asthma	71 (11.0)	33 (12.7)	18 (10.4)	20 (9.5)	.5434
CSU/CIU severity at the time of diagnosis (%)	642	259	173	210	.0101
Mild	71 (11.1)	27 (10.4)	32 (18.5)	12 (5.7)	
Moderate	202 (31.5)	81 (31.3)	53 (30.6)	68 (32.4)	
Severe	233 (36.3)	98 (37.8)	45 (26.0)	90 (42.9)	
Data not available	136 (21.2)	53 (20.5)	43 (24.9)	40 (19.0)	
UAS7 <sub>TD</sub>					
Mean (SD) UAS7 <sub>TD</sub>	17.3 (10.46)	17.6 (10.55)	14.6 (8.97)	19.2 (11.05)	.0032
UAS7 <sub>TD</sub> band (%)	600	244	161	195	.0140
0-6	98 (16.3)	41 (16.8)	28 (17.4)	29 (14.9)	
7-15	204 (34.0)	75 (30.7)	73 (45.3)	56 (28.7)	
16-27	182 (30.3)	82 (33.6)	40 (24.8)	60 (30.8)	
28-42	116 (19.3)	46 (18.9)	20 (12.4)	50 (25.6)	

CIU, chronic idiopathic urticaria; CSU, chronic spontaneous urticaria; Q1, first quartile; Q3, third quartile; MRA, medical record abstraction; NSAIDs, non-steroidal anti-inflammatory drugs; SD, standard deviation; UAS7<sub>TD</sub>, Urticaria Activity Score over 7 Days, twice-daily assessment.

<sup>a</sup>For categorical variables, the total number of patients with nonmissing data is presented by angioedema category for each question along with "n (%)" for each possible response to the question. Except where otherwise noted, percentage denominators are the number of patients with nonmissing data for the applicable question.

<sup>b</sup>P values shown from t tests for means of continuous variables and chi-square tests for frequencies of categorical variables to compare patients who experience angioedema (*Yes-angioedema*) with those who do not (*No-angioedema*). Patients in the *Misaligned* category were not included in the statistical comparison. For race and ethnicity, due to the small number of patients in many of the race/ethnicity categories, all races other than white were combined for these comparisons. Patients whose race/ethnicity group is "data not available" were excluded from the statistical comparison.

<sup>c</sup>Race and ethnicity data were not collected in France. For some categories of race and ethnicity, different terms were used across countries.


had severe (UAS7<sub>TD</sub> 28-42) disease activity (18.9% vs 12.4%) (Figure S1).<sup>19</sup> Moreover, a significant difference was observed between patients in the *Yes-angioedema* and *No-angioedema* categories in the mean (SD) absolute number of hives over a week collected in the UPDD (147.9 [266.37] vs 77.1 [123.87], respectively).

### 3.5 | Effect of angioedema on HRQoL, sleep and daily activities

The mean (SD) CU-Q<sub>2</sub>oL score for the overall patient sample was 33.6 (21.02). Mean (SD) CU-Q<sub>2</sub>oL scores differed significantly

between patients in the *Yes-angioedema* and *No-angioedema* categories (37.6 [20.81] vs 23.4 [17.12],  $P < .001$ ), reflecting significantly lower HRQoL for patients with angioedema (Figure 3A). Significant differences were observed between patients with and without angioedema on all CU-Q<sub>2</sub>oL domains.

On the DLQI, mean (SD) overall score for the entire sample was 9.1 (6.63) and for *Yes-angioedema* and *No-angioedema* groups were 10.4 (6.85) and 6.6 (5.21), respectively ( $P < .001$ ) (Figure 3B). Again, significant differences were observed between patients with and without angioedema on all DLQI domains (Figure 3C). More patients in the *Yes-angioedema* category (45.5%) than in the *No-angioedema*



**FIGURE 2** Mean urticaria activity score over 7 days, twice-daily assessment (UAS7<sub>TD</sub>), by angioedema classification. UAS7<sub>TD</sub> scores were calculated by summing the average of twice-daily assessments of hive count and itch score and summing these daily scores over 7 days. \* $P < .01$  (Yes—angioedema vs No—angioedema). Patients who fell into the *Misaligned* group were not included in the statistical comparisons. SD, standard deviation. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

category (20.4%) had DLQI scores  $>10$  (indicating a very to extremely large effect on HRQoL<sup>21</sup>).

As patients reported on the UPDD, mean (SD) weekly scores for interference with sleep were higher for patients in the *Yes-angioedema* category (6.9 [6.14]) than for patients in the *No-angioedema* category (4.6 [4.92]) ( $P < .001$ ), as were mean (SD) weekly scores for interference with daily activities (6.5 [5.68] vs 4.8 [4.39];  $P < .05$ ) (Figure 3D). Activity impairment measured by the mean percentage overall activity impairment score on the WPAI was also greater for patients in the *Yes-angioedema* category than for patients in the *No-angioedema* category (34.5% vs 23.8%;  $P < .001$ ) (Figure 4).

### 3.6 | Economic and societal impact of angioedema

Patients in the *Yes-angioedema* category had greater HCRU, both documented in medical records and self-reported, than patients in the *No-angioedema* category (Table S2). According to the MRA, patients in the *Yes-angioedema* category were more likely to have a CSU-related emergency department visit than patients in the *No-angioedema* category (24.7% vs 5.2%,  $P < .001$ ) and were more likely to have one or more inpatient hospital stays in the past 12 months (11.6% vs 3.5%,  $P < .01$ ). As reported in the patient diary, patients with angioedema were more likely to call a healthcare provider during the 7-day window than patients without angioedema (7.3% vs 0.6%,  $P < .01$ ).

Impact on work was greater among patients in the *Yes-angioedema* category than among patients in the *No-angioedema* category. Patients in the *Yes-angioedema* category who completed the diary

were more likely to have missed 1 or more hours of work in the 7-day diary window than patients in the *No-angioedema* category (27.6% vs 5.8%,  $P < .001$ ), and mean (SD) number of days of work missed in the past 3 months was significantly higher for patients in the *Yes-angioedema* versus the *No-angioedema* category (4.7 [11.64] vs 0.8 [3.86],  $P < .001$ ). WPAI scores showed significantly greater absenteeism among patients in the *Yes-angioedema* versus *No-angioedema* categories (mean [SD] percentage absenteeism: 9.1% [23.22%] vs 1.4% [9.08%],  $P < .001$ ), as well as significantly greater overall work impairment (mean [SD] percentage overall work impairment: 29.2% [28.48%] vs 19.1% [21.37%],  $P = .02$ ) (Figure 4). Overall, angioedema was the second most common reason for missing work, after itching.

### 3.7 | Characteristics of the misaligned angioedema category

Patients in the *Misaligned* category were compared descriptively with patients in the *Yes-angioedema* and *No-angioedema* categories and were found to be most similar to patients in the *Yes-angioedema* category in all outcomes. Demographics were similar in terms of age, sex, disease duration from diagnosis to enrolment and CSU severity at diagnosis (Table 1). Similar proportions of patients in the *Misaligned* and *Yes-angioedema* categories had moderate (UAS7<sub>TD</sub> 16-27) or severe (UAS7<sub>TD</sub> 28-42) disease activity (56.4% and 52.5%, respectively). When completing the survey, 52.2% of patients in the *Misaligned* category reported that their angioedema typically lasts more than 12 hours, compared with 47.8% of patients in the *Yes-angioedema* category. The mean (SD) amount of swelling, on a scale of 0 to 10, was similar for patients in the *Misaligned* category (7.1 [2.12]) and those in the *Yes-angioedema* category (7.2 [2.20]). In addition, the HRQoL impact of CSU was similar for patients in the *Misaligned* and *Yes-angioedema* categories, as indicated by mean (SD) overall scores on the DLQI (10.4 [6.85] and 9.7 [6.87], respectively) and CU-Q<sub>2</sub>oL (37.6 [20.81] and 37.0 [21.49], respectively) (Figure 3A,B). Percentage overall work impairment due to CSU was similar for patients in the *Misaligned* and *Yes-angioedema* categories (mean [SD] = 31.5% [29.99%] and 29.2% [28.48%], respectively) (Figure 4).

### 3.8 | DLQI regression analysis

Given the similarities in characteristics and outcomes among patients in the *Misaligned* and *Yes-angioedema* categories, the *Misaligned* patients were combined with the *Yes-angioedema* patients in the primary regression analysis of DLQI total score. After covariate adjustment, mean DLQI score was significantly higher for patients with angioedema versus without (9.88 vs 7.27,  $P < .001$ ) (Table 2). Results were similar in the sensitivity analysis in which patients in the *Misaligned* angioedema category were excluded: mean DLQI total score was significantly higher for patients with angioedema versus without (9.69 vs 6.73,  $P < .001$ ).

## 4 | DISCUSSION

The results of this post hoc analysis revealed notable differences in reporting of angioedema between CSU patients and physicians in the ASSURE-CSU study. The objective of this analysis was to align angioedema data in order to better estimate the actual rate of angioedema and to analyse differences between patients with and without angioedema. Angioedema was more frequently reported by patients than by physicians: 65.8% of patients who completed the survey reported having experienced angioedema during the past 12 months, whereas physicians reported that 41.0% of patients had experienced angioedema during the same time period. A large majority (97%) of patients with misaligned angioedema data did not have physician-reported angioedema but reported having experienced angioedema in either the patient survey or diary. These findings suggest that the occurrence of angioedema may be under-recognized among physicians (who may not routinely ask patients about angioedema symptoms), that patients may not always report their angioedema symptoms to physicians or that patients characterize some CSU-related swelling as angioedema when a physician would classify it as hives or sometimes as swelling unrelated to urticaria.

ASSURE-CSU patients with misaligned angioedema data appear most similar to patients with angioedema, in terms of disease duration; disease activity; and scores on patient-reported outcome measures including the CU-Q<sub>2oL</sub>, DLQI and WPAI. If patients in the *Misaligned* category were reclassified into the *Yes-angioedema* category, which they closely resembled, the percentage of patients in the ASSURE-CSU study with angioedema over 12 months would increase from 40.3% to 73.1%. Moreover, our analysis suggests that patients with more severe CSU are more likely to have angioedema. Physicians reported that more patients with angioedema than without had severe CSU at diagnosis, and patients with angioedema reported greater CSU activity as indicated by UAS7<sub>TD</sub> mean score than those without angioedema.

The primary ASSURE-CSU analysis found that angioedema had a significant impact on HRQoL in patients with CSU, particularly with respect to emotional well-being, fatigue and mood.<sup>12</sup> The current analysis also revealed the considerable impact of angioedema on HRQoL, as indicated by CU-Q<sub>2oL</sub> and DLQI scores, and its additional negative impact on top of itch and hives. Even after controlling for other factors, angioedema has a significant effect on the DLQI total score. In addition, patients with angioedema have additional impairment on work productivity and more HCRU than those without it, indicating that angioedema further contributes to the economic burden of inadequately controlled CSU.

Taken together, findings from these analyses suggest that physicians should ensure that patients understand the manifestations of angioedema and should routinely assess patients with CSU for presence of angioedema, which may enable more accurate estimates of the true prevalence and burden of angioedema. Appropriate tools available for evaluating angioedema activity and its impact on CSU patients' lives should be used regularly in clinical practice.<sup>22-24</sup> In patients with recurrent angioedema, CSU should be considered as one of the differential diagnoses.<sup>1,25</sup> Additionally, angioedema in CSU should prompt adequate treatment. The development of new treatments specifically for CSU has been shown to prevent angioedema and improve angioedema-dependent and dermatology-related quality of life.<sup>9,26-28</sup> Future research should explore the clinical course of angioedema in CSU—including the most commonly affected areas, the clinical differences compared with hereditary angioedema and evolution of symptoms over time—as well as the distinct management patterns that patients with angioedema in CSU require. Understanding the underlying mechanisms of angioedema in CSU<sup>29</sup> and exploring differences in the general categories of angioedema (ie mast cell-induced and bradykinin-induced) are also important areas for additional research.


Several limitations must be considered when the results of this study are interpreted. First, the study was conducted in specialized centres only, and physicians and centres were not selected systematically, owing to the small number of specialist units for this population. The resulting sample is not guaranteed to be representative of all medical settings and physicians treating patients with CSU. Second, the target population had had CSU for at least 12 months and had not responded adequately to treatment at the time of inclusion in the study; thus, the study does not reflect the entire population of patients with CSU but more patients with inadequately controlled CSU. Moreover, although patients received training about identifying angioedema and reporting it in the patient survey or diary, like all patient-reported data, angioedema was still subject to patients' individual interpretation of the symptom description and manifestation. In addition, data extracted by physicians from medical records were not independently validated.

## 5 | CONCLUSIONS


Among this study population with inadequately controlled CSU, 40.3% had confirmed angioedema in the past 12 months, and 32.8%

**FIGURE 3** Impact of CSU on HRQoL, sleep and daily activities, overall and by angioedema classification. (A) CU-Q<sub>2oL</sub> total and domain scores. Different CU-Q<sub>2oL</sub> scale scores are used in Germany than in Canada, France, Italy, the Netherlands, Spain and the UK. German CU-Q<sub>2oL</sub> overall score and scale scores are presented in Figure S2; (B) DLQI total score; (C) DLQI domain scores; (D) Interference with sleep and interference with daily activities over a week, as reported on the Urticaria Patient Daily Diary, by angioedema classification. For all scores, a higher score means higher impairment. \**P* < .001 (Yes—angioedema vs No—angioedema); \*\**P* < .05 (Yes—angioedema vs No—angioedema). Patients who fell into the *misaligned* category were not included in the statistical comparisons. CU-Q<sub>2oL</sub>, Chronic Urticaria Quality of Life Questionnaire; DLQI, Dermatology Life Quality Index; HRQoL, health-related quality of life; SD, standard deviation. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]


(A) CU-Q<sub>2</sub>oL Total and Domain Scores




(B) DLQI total score\*




(C) DLQI domain scores



(D) Severe interference





**FIGURE 4** Work productivity and activity impairment (WPAI) results by angioedema classification. Absenteeism was defined as percentage of work time missed due to chronic spontaneous urticaria (CSU) in the past 7 days. Presenteeism was defined as percentage impairment while working due to CSU in the past 7 days. Overall work impairment was defined as percentage work impairment due to CSU in the past 7 days, incorporating both absenteeism and presenteeism using the following validated WPAI algorithm: overall work impairment = absenteeism + (1-absenteeism)\*presenteeism. \* $P < .001$  (Yes—angioedema vs No—angioedema); \*\* $P < .05$  (Yes—angioedema vs No—angioedema). Patients who fell into the misaligned category were not included in the statistical comparisons. SD, standard deviation; WPAI, Work Productivity and Activity Impairment. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

**TABLE 2** DLQI regression results

Angioedema Classification	Within Group			Contrast in LS Mean				
	n	LS Mean	SE	Comparison	LS Mean	SE	95% CI	P value
Primary analysis: ANCOVA for DLQI total score by angioedema classification (assigning <i>Misaligned</i> category to <i>Yes-angioedema</i> category)								
No-angioedema	150	7.27	0.850	vs Yes	-2.61	0.558	-3.71 to -1.51	<.0001
Yes-angioedema	399	9.88	0.722					
Sensitivity analysis: ANCOVA for DLQI total score by angioedema classification (removing the <i>Misaligned</i> category)								
No-angioedema	150	6.73	0.902	vs Yes	-2.96	0.586	-4.12 to -1.81	<.0001
Yes-angioedema	232	9.69	0.795					

LS means, 95% CIs and  $P$  values are from an ANCOVA model with covariates: angioedema classification, UAS7<sub>TD</sub> score (continuous), age at enrolment (continuous), sex (male and female), country (Canada, France, Germany, Italy, Spain, the Netherlands and the UK), disease duration from diagnosis to enrolment (continuous) and comorbidities at enrolment (hypersensitivity to NSAIDs [yes/no], Hashimoto's [yes/no] and asthma [yes/no]). ANCOVA, analysis of covariance; CI, confidence interval; DLQI, Dermatology Life Quality Index; LS, least squares; n, number of subjects with data for all model inputs; NSAIDs, nonsteroidal anti-inflammatory drugs; SE, standard error; UAS7<sub>TD</sub>, Urticaria Activity Score over 7 days, twice-daily assessment.

of patients had misaligned angioedema data (ie the majority reporting having experienced angioedema in the past 12 months without having it documented in their medical records). This suggests that a higher proportion of patients with inadequately controlled symptomatic CSU might have angioedema than previously thought. Patients with angioedema reported statistically significantly worse HRQoL and higher societal burden than patients without angioedema; patients in the Misaligned group reported similar impairment as patients with confirmed angioedema. Overall, the study found that angioedema has an incremental impact on societal and humanistic

outcomes in CSU patients. These findings suggest the need for improved physician-patient communication regarding angioedema for better symptom control in patients with inadequately controlled CSU.

#### ACKNOWLEDGEMENTS

The authors gratefully acknowledge the following recruiting physicians: Drs. Sameh Hanna, Jacques Hebert, Amin Kanani, Paul Keith, Gina Lacuesta, Jason Lee, G-Daniel Schachter, Susan Wasserman and Shahin Zanganeh in Canada; Drs. Emmanuelle Amsler-Soria, Annick

Barbaud, Claire Bernier, Laurence Bouillet, Jean-Jacques Grob, Giordano Labadie, Laurent Machet, Paul Martin, Fabien Pelletier, Nadia Rasion-Peyron, Delphine Staumont-Salle and Manuelle Viguier in France; Drs. Andrea Bauer, Randolph Brehler, Hans Merk, Franziska Rueff, Petra Staubach-Renz and Amir Yazdi in Germany; Drs. Ornella de Pità, Silvia Mariel Ferrucci, Maria Laura Flori, Giampiero Girolomoni, Giovanni Pellacani, Paolo Pigatto and Domenico Schiavino in Italy; Drs. Menno T. W. Gaastra, G. R. R. Kuiters, M. L. A. Schutte-laar, R. A. Tupker, Thomas Rustemeyer, Phyllis Spuls and Roy Gerth van Wijk in the Netherlands; Drs. Jesús Borbujo, Pablo de la Cueva, Alejandro Joral Badas, Moisés Labrador Hornillo, Ana Perez Montero, Javier Pedraz and Esther Serra in Spain; and Drs. Anthony Bewley, Seatak Cheung, Nyz Chiang, Venkata Gudi, Frances Humphreys, Dimtra Koumaki, John Reed and Donna Torley in the United Kingdom.

Kate Lothman of RTI Health Solutions provided medical writing services, which were funded by Novartis Pharma AG.

### CONFLICT OF INTEREST

This research was performed under a research contract between RTI Health Solutions and Novartis Pharma AG and was funded by Novartis Pharma AG. K.H., S.H., D.McB., C.S. and D.W. are employees of RTI Health Solutions, which provides consulting and other research services to pharmaceutical, device, governmental and non-government organizations. In their salaried positions, they work with a variety of companies and organizations. They receive no payment or honoraria directly from these organizations for services rendered. M.M.B. is an employee of Novartis Pharma AG. H.T. is an employee of Novartis Pharmaceuticals Corporation.

### AUTHOR CONTRIBUTIONS

M.M.B., K.H. and D.McB. initiated the study. M.M.B., K.H., D.McB., C.S. and D.W. designed the study; M.A., F.B., W.C., H.O.E., A.G-A., C.G., A.K., J-P.L., C.L., A.M., M.M., A.N., J.O.d.F., G.S. and K.W. provided clinical input on the study design. K.H., D.McB., C.S. and D.W. managed data collection, and M.M.B., K.H., D.McB., S.H., C.S., H.T. and D.W. led the data analyses. M.M.B., M.A., F.B., W.C., H.O.E., A.G-A., C.G., K.H., A.K., J-P.L., C.L., A.M., M.M., D.McB., A.N., J.O.d.F., C.P., G.S., C.S., H.T., K.W. and D.W. interpreted the data. M.M.B., K.H., S.H., D.McB. and G.S. drafted the manuscript, and M.A., F.B., W.C., H.O.E., A.G-A., C.G., A.K., J-P.L., C.L., A.M., M.M., A.N., J.O.d.F., C.S., H.T., K.W. and D.W. revised it critically for intellectual content. All authors reviewed and approved the final manuscript and agree to be accountable for the work as a whole.

### ORCID

G. Sussman  <http://orcid.org/0000-0002-2202-2513>  
 M. Abuzakouk  <http://orcid.org/0000-0003-0802-8342>  
 W. Canonica  <http://orcid.org/0000-0001-8467-2557>  
 A. Giménez-Arnau  <http://orcid.org/0000-0001-5434-7753>

C. Grattan  <http://orcid.org/0000-0002-8466-4351>  
 A. Knulst  <http://orcid.org/0000-0002-1056-3179>  
 J.-P. Lacour  <http://orcid.org/0000-0001-7663-2053>  
 D. McBride  <http://orcid.org/0000-0002-2545-651X>  
 M. Maurer  <http://orcid.org/0000-0002-4121-481X>  
 A. Nakonechna  <http://orcid.org/0000-0002-0141-6361>  
 K. Weller  <http://orcid.org/0000-0003-4437-0313>  
 M.-M. Balp  <http://orcid.org/0000-0002-8612-7680>

### REFERENCES

- Zuberbier T, Aberer W, Asero R, et al. The EAACI/GA<sup>2</sup>LEN/EDF/WAO guideline for the definition, classification, diagnosis, and management of urticaria: the 2013 revision and update. *Allergy*. 2014;69:868-887.
- Rye Rasmussen EH, Bindslev-Jensen C, Bygum A. Angioedema—assessment and treatment. *Tidsskr Nor Laegeforen*. 2012;132:2391-2395.
- Banerji A, Sheffer AL. The spectrum of chronic angioedema. *Allergy Asthma Proc*. 2009;30:11-16.
- Maurer M, Weller K, Bindslev-Jensen C, et al. Unmet clinical needs in chronic spontaneous urticaria: a GA<sup>2</sup>LEN task force report. *Allergy*. 2011;66:317-330.
- Maurer M, Staubach P, Raap U, et al. H1-antihistamine-refractory chronic spontaneous urticaria: it's worse than we thought—first results of the multicenter real-life AWARE study. *Clin Exp Allergy*. 2017a;47:684-692.
- van der Valk P, Moret G, Kiemeny L. The natural history of chronic urticaria and angioedema in patients visiting a tertiary referral centre. *Br J Dermatol*. 2002;146:110-113.
- Sánchez-Borges M, Caballero-Fonseca F, Capriles-Hulett A, González-Aveledo L, Maurer M. Factors linked to disease severity and time to remission in patients with chronic spontaneous urticaria. *J Eur Acad Dermatol Venereol*. 2017;31:964-971.
- Weller K, Groffik A, Magerl M, et al. Development and construct validation of the angioedema quality of life questionnaire. *Allergy*. 2012;67:1289-1298.
- Staubach P, Metz M, Chapman-Rothe N, et al. Effect of omalizumab on angioedema in H1-antihistamine resistant chronic spontaneous urticaria patients: results from X-ACT, a randomized controlled trial. *Allergy*. 2016;71:1135-1144.
- Lumry WR, Castaldo AJ, Vernon MK, Blaustein MB, Wilson DA, Horn PT. The humanistic burden of hereditary angioedema: impact on health-related quality of life, productivity, and depression. *Allergy Asthma Proc*. 2010;31:407-414.
- Weller K, Maurer M, Grattan C, et al. ASSURE-CSU: a real-world study of burden of disease in patients with symptomatic chronic spontaneous urticaria. *Clin Transl Allergy*. 2015a;5:29.
- Maurer M, Abuzakouk M, Bérard F, et al. The burden of chronic spontaneous urticaria is substantial: real-world evidence from ASSURE-CSU. *Allergy*. 2017b;72:2005-2016.
- World Medical Association. Declaration of Helsinki: ethical principles for medical research involving human subjects. October 2008. Available at: <https://www.wma.net/policies-post/wma-declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/>. Accessed April 16, 2016.
- Baiardini I, Pasquali M, Braidò F, et al. A new tool to evaluate the impact of chronic urticaria on quality of life: chronic urticaria quality of life questionnaire (CU-Q2oL). *Allergy*. 2005;60:1073-1078.
- Młynek A, Magerl M, Hanna M, et al. The German version of the Chronic Urticaria Quality-of-Life Questionnaire: factor analysis, validation, and initial clinical findings. *Allergy*. 2009;64:927-936.

16. Finlay AY, Khan GK. Dermatology Life Quality Index (DLQI)—a simple practical measure for routine clinical use. *Clin Exp Dermatol.* 1994;19:210-216.
17. Flood EM, Zazzali JL, Devlen J. Demonstrating measurement equivalence of the electronic and paper formats of the urticaria patient daily diary in patients with chronic idiopathic urticaria. *Patient.* 2013;6:225-231.
18. Mathias SD, Dreskin SC, Kaplan A, Saini SS, Spector S, Rosén KE. Development of a daily diary for patients with chronic idiopathic urticaria. *Ann Allergy Asthma Immunol.* 2010;105:142-148.
19. Reilly M. Work Productivity and Activity Impairment Questionnaire. 2004. <http://www.reillyassociates.net/Index.html>. Accessed December 14, 2013.
20. Stull D, McBride D, Tian H, et al. Analysis of disease activity categories in chronic spontaneous/idiopathic urticaria. *Br J Dermatol.* 2017;177:1093-1101.
21. Hongbo Y, Thomas CL, Harrison MA, Salek MS, Finlay AY. Translating the science of quality of life into practice: what do dermatology life quality index scores mean? *J Invest Dermatol.* 2005;125:659-664.
22. Weller K, Zuberbier T, Maurer M. Clinically relevant outcome measures for assessing disease activity, disease control and quality of life impairment in patients with chronic spontaneous urticaria and recurrent angioedema. *Curr Opin Allergy Clin Immunol.* 2015b;15:220-226.
23. Weller K, Magerl M, Peveling-Oberhag A, Martus P, Staubach P, Maurer M. The angioedema quality of life questionnaire (AE-QoL) – assessment of sensitivity to change and minimal clinically important difference. *Allergy.* 2016;71:1203-1209.
24. Weller K, Groffik A, Magerl M, et al. Development, validation, and initial results of the angioedema activity score. *Allergy.* 2013;68:1185-1192.
25. Maurer M, Magerl M, Metz M, Siebenhaar F, Weller K, Krause K. Practical algorithm for diagnosing patients with recurrent wheals or angioedema. *Allergy.* 2013;68:816-819.
26. Maurer M, Sofen H, Ortiz B, Kianifard F, Gabriel S, Bernstein JA. Positive impact of omalizumab on angioedema and quality of life in patients with refractory chronic idiopathic/spontaneous urticaria: analyses according to the presence or absence of angioedema. *J Eur Acad Dermatol Venereol.* 2017c;31:1056-1063.
27. Zazzali JL, Kaplan A, Maurer M, et al. Angioedema in the omalizumab chronic idiopathic/spontaneous urticaria pivotal studies. *Ann Allergy Asthma Immunol.* 2016;117:370-377.
28. Finlay AY, Kaplan AP, Beck LA, et al. Omalizumab substantially improves dermatology-related quality of life in patients with chronic spontaneous urticaria. *J Eur Acad Dermatol Venereol.* 2017;31:1715-1721.
29. Terhorst D, Koti I, Krause K, Metz M, Maurer M. In chronic spontaneous urticaria, high numbers of dermal endothelial cells, but not mast cells, are linked to recurrent angioedema. *Clin Exp Dermatol.* 2017;43:131-136.

## SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

**How to cite this article:** Sussman G, Abuzakouk M, Bérard F, et al. Angioedema in chronic spontaneous urticaria is underdiagnosed and has a substantial impact: Analyses from ASSURE-CSU. *Allergy.* 2018;73:1724–1734. <https://doi.org/10.1111/all.13430>